



Williamson, W., Lewandowski, A. J., Forkert, N. D., Griffanti, L., Okell, T. W., Betts, J., Boardman, H., Siepmann, T., McKean, D., Huckstep, O., Francis, J. M., Neubauer, S., Phellan, R., Jenkinson, M., Doherty, A., Dawes, H., Frangou, E., Malamateniou, C., Foster, C., & Leeson, P. (2018). Association of cardiovascular risk factors with MRI indices of cerebrovascular structure and function and white matter hyperintensities in young adults. *JAMA - Journal of the American Medical Association*, 320(7), 665-673.
<https://doi.org/10.1001/jama.2018.11498>

Peer reviewed version

Link to published version (if available):
[10.1001/jama.2018.11498](https://doi.org/10.1001/jama.2018.11498)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via AMA at <https://jamanetwork.com/journals/jama/fullarticle/2697697> . Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

1 ***Association of cardiovascular risk factors with MRI indices of cerebrovascular***
2 ***structure and function and white matter hyperintensities in young adults***

3 Wilby Williamson¹, MSc, MRCP, Adam J Lewandowski^{1,2}, DPhil, Nils D
4 Forkert³, PhD, Ludo Griffanti⁴, PhD, Thomas W Okell⁴, DPhil, Jill Betts¹, DPhil,
5 Henry Boardman¹, MRCP, DPhil, Timo Siepmann⁵ MD, PhD, David McKean⁶,
6 MD, Odaro Huckstep¹, MSc, Jane Francis², DCR(R), Stefan Neubauer², MD,
7 FRCP, DPhil, Renzo Phellan³, MSc Mark Jenkinson⁴, DPhil, Aiden Doherty⁷,
8 PhD, Helen Dawes⁸, PhD, Eleni Frangou⁹, MSc (Res), Christina
9 Malamateniou^{10,11}, PhD, Charlie Foster¹², PhD, Paul Leeson¹, PhD, FRCP*

10 ¹ Oxford Cardiovascular Clinical Research Facility and ² Oxford Centre for
11 Clinical Magnetic Resonance Research, Division of Cardiovascular Medicine,
12 Radcliffe Department of Medicine, University of Oxford, Oxford, UK.

13 ³ Department of Radiology and Hotchkiss Brain Institute, University of Calgary,
14 Calgary, Alberta, Canada

15 ⁴ Wellcome Centre for Integrative Neuroimaging, FMRIB Division, Nuffield
16 Department of Clinical Neurosciences, University of Oxford, Oxford, UK

17 ⁵ Department of Neurology, University Hospital Carl Gustav Carus, Technische
18 Universität Dresden, Dresden, Germany.

19 ⁶ Department of Radiology, Stoke Mandeville Hospital, Buckinghamshire NHS
20 Trust, UK

21 ⁷ Nuffield Department of Population Health, BHF Centre of Research
22 Excellence and Big Data Institute, Li Ka Shing Centre for Health Information
23 and Discovery, University of Oxford

24 ⁸ Faculty of Health and Life Sciences, Oxford Brookes University, Oxford

25 ⁹ Centre for Statistics in Medicine, Nuffield Department of Orthopaedics,
26 Rheumatology and Musculoskeletal Sciences, University of Oxford, UK

27 ¹⁰ Imaging and Biomedical Engineering Clinical Academic Group, Kings College
28 London and ¹¹ Department of Family Care and Mental Health, University of
29 Greenwich UK

30 ¹² School of Policy Studies, University of Bristol, Bristol, UK

31 *Correspondence to: Professor Paul Leeson, Oxford Cardiovascular Clinical
32 Research Facility, Division of Cardiovascular Medicine, Radcliffe Department of
33 Medicine, University of Oxford, John Radcliffe Hospital, Oxford. OX3 9DU, UK.

34 Tel: +44 1865 572846. Fax +44 1865 221111. E-mail:

35 paul.leeson@cardiov.ox.ac.uk

36 **Date of Revision 11th July 2018**

37 **Word Count 3105**

38 **Key Points**

39 **Questions:** Are modifiable cardiovascular risk factors in young adults
40 associated with cerebral blood vessel structure and function, and neuroimaging
41 white matter hyperintensities?

42 **Results:** In this cross-sectional study of 125 young adults without clinical
43 evidence of cerebrovascular disease, a higher number of optimal
44 cardiovascular health metrics was correlated with higher cerebral vessel
45 density, higher cerebral blood flow, and lower white matter hyperintensity
46 lesions.

47 **Meaning:** These preliminary findings suggest a relationship between
48 modifiable cardiovascular risk factors and MRI biomarkers of cerebrovascular
49 structure and function and white matter hyperintensities in young adults.
50 Further research is needed to verify these findings and determine clinical
51 importance.

52

53 **Abstract**

54 **Importance:** Risk of stroke and brain atrophy in later life relate to levels of
55 cardiovascular risk in early adulthood. However, it is unknown whether
56 cerebrovascular changes are present in young adults.

57 **Objective:** To examine relationships between modifiable cardiovascular risk
58 factors and cerebrovascular structure, function and white matter integrity in
59 young adults.

60 **Design, Setting, and Participants:** A cross-sectional observational study of
61 125 young adults (aged 18 to 40 years) without clinical evidence of
62 cerebrovascular disease with data collection completed between August 2014
63 and May 2016 at the University of Oxford, United Kingdom. Final data
64 collection was completed on the 31st of May 2016.

65 **Exposures:** The number of modifiable cardiovascular risk factors at
66 recommended levels, based on the following criteria: BMI <25 kg/m²; highest
67 tertile cardiovascular fitness and/or physical activity; alcohol <8 drinks/week;
68 non-smoker for >6 months; blood pressure on awake ambulatory monitoring
69 <130/80 mmHg; a non-hypertensive diastolic response to exercise (peak
70 diastolic blood pressure <90 mmHg), total cholesterol <200mg/dL; and fasting
71 glucose <100mg/dL. Participants were categorized from 0 to 8, with higher
72 numbers indicating healthier risk categories.

73 **Main Outcomes and Measures:** Cerebral vessel density (vessels/cm³), caliber
74 (μm) and tortuosity, brain white matter hyperintensity lesion count (number),
75 and in a subgroup (n=52) brain blood arrival time (seconds) and cerebral blood
76 flow (ml/100g/min) assessed by brain magnetic resonance imaging.

Results A total of 125 participants, mean age 25±5 years, 49% female, with a mean score of 6.0 (SD 1.4) modifiable cardiovascular risk factors at recommended levels, completed the cardiovascular risk assessment and brain MRI protocol.

Cardiovascular risk factors were correlated with cerebrovascular morphology and white matter hyperintensity count in multivariable models. For each additional modifiable risk factor categorized as healthy, vessel density was greater by 0.3 vessels/cm³ (95%CI 0.1 to 0.5, p=0.003), vessel caliber was greater by 8µm (95%CI 3 to 13, p=0.01) and white matter hyperintensity lesions was lower by 1.6 lesions (95%CI 0.5 to 2.8, p=0.006). Among the 52 participants with available data, cerebral blood flow varied with vessel density and was 2.5ml/100g/min higher for each healthier category of a modifiable risk factor (95%CI 0.16 to 4.89, p=0.03).

Conclusions and Relevance In this preliminary study, involving young adults without clinical evidence of cerebrovascular disease, modifiable cardiovascular risk factors were associated with MR indices of cerebral vessel structure and function, and white matter hyperintensities. Further research is needed to verify these findings and determine clinical importance.

Key words: brain health, cardiovascular risk factors, young adults,

Introduction

A life-course approach to understand risk of cardiovascular disease is well established¹ and it is accepted that changes in cardiac and vascular structure that underlie this risk emerge very early in life^{2, 3}. Whether modifiable cardiovascular risk factors, and novel early life exposures such as preterm birth, influence the early cerebrovasculature is less well studied.

Cardiovascular risk closely relates to cerebrovascular injury and cognitive decline in older adults⁴. MRI markers of cerebral injury in mid-life, including white matter hyperintensity lesions, are associated with future stroke, dementia and all-cause mortality⁵. Progression of white matter hyperintensity lesions is faster in association with metabolic dysfunction and hypertension⁶. Experimental studies have demonstrated that cardiovascular risk factors result in remodelling of the brain vasculature, including vessel rarefaction, lower vessel caliber and cerebral blood flow⁷. Elevated blood pressure, dyslipidemia and low fitness in early adulthood are known to predict brain health in older adult life⁸⁻¹⁰. Whether cerebrovascular morphological changes are already evident in young adults, and correlate with white matter hyperintensity lesions and risk factors at this age, is unclear.

Advances in brain MRI allow automated segmentation and analysis of vessel morphology, white matter hyperintensity lesions^{11, 12} and blood flow¹³; thus making it possible to estimate brain vascular and structural status for an individual^{11, 12}. Therefore, the objective of the current study was to use multi-modality brain imaging to test the hypothesis that cardiovascular risk profiles are correlated with variation in vessel morphology and white matter hyperintensity lesions in young adults.

Methods

Study design and participants

This was a cross-sectional observational study completed between August 2014 and May 2016. The South Central Research Ethics Committee for the National Health Service Health Research Authority (NHS HRA) approved the study (14/SC/0275). All participants gave written informed consent. Measurements were completed at the Oxford Cardiovascular Clinical Research Facility and Oxford Centre of Clinical Magnetic Resonance Research, John Radcliffe Hospital, University of Oxford, United Kingdom. Image analysis was performed using pipelines developed at the Hotchkiss Brain Institute, University of Calgary and Wellcome Centre for Integrative Neuroimaging, University of Oxford¹²⁻¹⁶. Final data collection was completed on the 31st of May 2016.

Participants aged 18 to 40 years were recruited through purposive active and passive recruitment¹⁷ including local advertising, invitation from local birth cohort studies and invitation from the Oxford University Hospital Hypertension Service. Strategies were designed to recruit adults with a heterogeneity in risk factors known to be present in young adult populations including traditional risk factors such as hypertension and more novel factors such as gestational age. Participants were excluded if they had previous cardiovascular or cerebrovascular events, renal dysfunction or metabolic disease including diagnosis of familial hyperlipidaemia. Participants with secondary causes of hypertension such as renal vascular disease, vascular anomalies or adrenal dysfunction were excluded following assessment in Oxford University Hospital Hypertension Service.

Procedures

Cardiovascular Risk Assessment

Participants attended a research clinic in the morning after a 12-hour fast to complete a detailed cardiovascular risk assessment (Supplementary Data eMethods 1).

Measurements included: body size, fasting blood samples for total cholesterol, high-density lipoprotein, low-density lipoprotein, triglycerides, highly sensitive c-reactive protein, glucose, and insulin levels, clinic and 24-hour blood pressure, as well as peak oxygen uptake and exercise blood pressure (from cardiopulmonary exercise testing). In addition, participants completed a detailed lifestyle questionnaire and had seven complete days of objectively measured physical activity. Post-hoc, participants were assessed for a cardiovascular score based on 8 modifiable risk factors, with 1 point awarded for each healthier category of a modifiable risk factor according to the following criteria: BMI $<25 \text{ kg/m}^2$; highest tertile cardiovascular fitness and/or physical activity; alcohol <8 drinks/week; non-smoker for >6 months; blood pressure on awake ambulatory monitoring $<130/80 \text{ mmHg}$; a non-hypertensive diastolic response to exercise (peak diastolic blood pressure $<90 \text{ mmHg}$), total cholesterol $<200 \text{ mg/dL}$; and fasting glucose $<100 \text{ mg/dL}$. The score was adapted from established cardiovascular health scores to include alcohol consumption and dynamic exercise blood pressure response, as known independent risk factors for brain health¹⁹⁻²¹. The thresholds for healthy criteria were set to be consistent with recommended public health guidelines and existing literature^{4, 9, 18-20}.

Brain Imaging and Analysis

Individuals underwent a multimodality MRI scan (3.0T Trio Tim, Siemens, Munich, Germany). The MRI protocol included T1-weighted structural, T2-weighted Fluid-Attenuated Inversion Recovery (FLAIR), Diffusion Tensor Imaging (DTI) and Time-of-Flight (TOF) MR Arteriogram (MRA) (Supplementary Data eMethods 2). MR imaging

was completed fasted and prior to exercise testing. Complete acquisition and analysis methods are presented in the on-line supplement.

T1-weighted images were processed using FMRIB Software Library (FSL) tools²¹. Brain vessel segmentation was completed on TOF MRA using previously described automated segmentation tools (supplement eFigure 1)^{12, 16}. Binary segmentations were used to determine vessel density, caliber and tortuosity.

White matter hyperintensity (WMH) lesions and associated volumes were segmented using the Brain Intensity AbNormality Classification Algorithm (BIANCA); a fully-automated, supervised method for WMH detection^{11, 22}. BIANCA classifies image voxels based on their intensity and spatial features, where the intensity features were extracted from T2-weighted FLAIR, T1-weighted and DTI fractional anisotropy (FA) images, FA images were generated using DTI tools, FSL topup, FSL eddy and DTIFit^{21, 23}. WMH masks were manually segmented from 10 images to use as the training set for BIANCA, these were independently verified by a neurologist (TS) and radiologist (DM) blinded to participant risk profile. Lesion count was selected as the most sensitive outcome of white matter change in young adults in whom a single lesion, independent of volume, could be considered abnormal²⁴. Minimum lesion size used in analysis was 1 mm³.

A subgroup of 52 participants also had multi-delay vessel-encoded pseudocontinuous Arterial Spin Labelling (ASL), identical to a previously published protocol¹³. Cerebral blood flow and blood arrival time were estimated from ASL images using a previously described analysis pipeline^{13, 15}. Gray matter masks were used to calculate the average cerebral blood flow after linear registration of the ASL MRI to the T1-weighted MRI dataset.

Statistical Analysis

Recruitment was continued to 125 participants for an estimated power of 90% at $P=0.05$ to identify a 0.70-SD difference in vessel density, vessel caliber and white matter lesion count between lowest and highest cardiovascular risk tertile groups. Arterial spin labelling (ASL) MRI imaging was added during the course of the study to provide a subgroup of 52 participants, recruited sequentially for an estimated 80% power to detect a 10% difference in cerebral blood flow²⁵.

Existing literature on risk predictors of brain health was used to define an a priori set of potential correlates of MRI brain health in young adults^{4, 9, 18-20}. These were grouped as: 1) non-modifiable, including age, sex, gestational age, and 2) modifiable, including systolic blood pressure, body mass index (BMI), peak exercise capacity (oxygen uptake ml/min/kg), peak exercise diastolic blood pressure, weekly vigorous activity level, alcohol consumption, smoking history, lipid profile, glucose and insulin resistance, and current hypertension medication.

In a priori analysis, bivariable and multivariable analysis was completed to investigate correlation between the defined cardiovascular risk markers and brain imaging findings. In this multivariable analysis to reduce multiple testing and potential interaction between the variables, the model was restricted to a subset of variables (resting systolic blood pressure, BMI, vigorous physical activity level, alcohol consumption and smoking). This model was adjusted for non-modifiable factors including age, sex and gestational age.

In post-hoc analysis, the individuals' combined cardiovascular score from across 8 risk factors, was used as a metric of overall modifiable cardiovascular health. The relationships between the individuals' modifiable cardiovascular score and brain

imaging findings were studied using linear regression adjusted for age and sex.

Comparison between brain imaging findings was made between groups of participants in the lowest and highest tertiles for the cardiovascular score.

In addition, bivariable analysis was completed to investigate correlation between vessel morphology and white matter hyperintensity lesion count and in a subgroup (n=52), blood arrival time and cerebral blood flow. These relationships were further investigated with fixed entry linear regression models adjusted for modifiable and non-modifiable factors used in the models above.

Statistical analysis was undertaken using Statistical Product and Service Solutions (SPSS) Version 22 (Armonk, New York, U.S). Normality of variables was assessed by visual assessment of curves. If normally distributed, results are presented as mean \pm standard deviation for continuous variables, otherwise median and interquartile range. For categorical variables, number and percentage are presented. Comparison between groups for continuous variables was performed with a 2-sided, independent-sample Student's *t* test. All multivariable analysis was completed using forced entry linear regression with residual analysis completed to assess model assumptions. All multivariable analyses were adjusted for age and sex. All tests were 2-sided, P-values <0.05 were considered statistically significant with no adjustment for multiple comparison. Due to multiplicity of testing all results were considered exploratory. Results are presented as point estimates and 95% confidence intervals stated in units appropriate to the risk factor and brain imaging findings being reported. Graphpad Prism 7 software was used for statistical figures and mean with 95% confidence intervals presented.

Results

A total of 125 participants completed the brain MRI protocol and cardiovascular risk assessment study measures. The mean age of participants was 24.7 ± 5.0 years, 61 participants were female (49%), the mean gestational age was 36.6 ± 4.3 weeks, educational attainment was high with 86 completing University level education (68.8%), 29 participants had prior history of hypertension of which 21 were on anti-hypertension medications (16.8%) (Table 1). The distributions of MRI brain outcomes between lowest and highest quintile of the respective measures are presented in the supplement (eTable 1). The 52 participants with available cerebral blood flow data shared a comparable demographic profile as the overall study group (mean age 24.6 ± 5.0 years, 42% female ($n=22$), gestational age 37.2 ± 3.6 weeks, and 10 participants were on anti-hypertension medications (19.2%).

Modifiable risk factors and association with brain vessel structure and white matter hyperintensity lesions

Association between risk factors (SBP, BMI, smoking pack years, Ex DBP, cholesterol/HDL ratio, hypertension treatment) and brain vessel morphology are presented in Table 2. Vessel tortuosity only varied with gestational age in both bivariable and adjusted models (0.005 unit tortuosity change/gestational week, 95%CI 0.001 to 0.009, $p=0.007$) (Table 2 and Supplementary Data, eTable 2). In the multivariable models, systolic blood pressure (-0.2 vessels/ cm^3 per 10mmHg, 95%CI -0.4 to -0.004 , $p=0.04$), smoking (2 vessels/ cm^3 per 10 pack years, 95%CI 0.6 to 3.0 , $p=0.04$) and body mass index (-0.1 vessels/ cm^3 per $1\text{kg}/\text{m}^2$, 95%CI -0.15 to -0.01 , $p=0.02$) were significantly correlated with vessel density, while vessel caliber was correlated with systolic blood pressure ($-6\mu\text{m}$ per 10mmHg, 95%CI -10.0 to -0.5 , $p=0.03$) and smoking ($40\mu\text{m}$ per 10 pack years, 95%CI 2.0 to 80.0 , $p=0.04$). In bivariable models, number of white matter hyperintensity lesions correlated with

smoking (8 lesions per 10 pack years, 95%CI 1.5 to 14.4, $p=0.02$), exercise diastolic blood pressure (1 lesions per 10mmHg, 95%CI 0.1 to 2.4, $p=0.04$), and alcohol consumption (4 lesion per 10 weekly alcoholic drinks, 95%CI 0.3 to 8.0, $p=0.03$), (Supplementary Data, eTable 3).

Healthier categories on the modifiable cardiovascular score correlated with vessel morphology (Table 4.) Each additional healthier category of risk factor was associated with a 0.3 vessels/cm³ higher vessel density (95%CI 0.1 to 0.5, $p=0.003$) and 8µm greater vessel caliber (95%CI 3.0 to 13.0, $p=0.01$). Similarly, white matter hyperintensity lesion count correlated with the cardiovascular score with 1.6 fewer white matter hyperintensity lesions per additional healthier category of risk factor (95%CI to -3.0 to -0.5, $p=0.006$). In addition, the cardiovascular score correlated with total volume of white matter hyperintensity adjusted for brain size with 51 mm³ lower white matter hyperintensity lesion volume per additional healthier category of risk factor (95%CI to -87 to -15 mm³ $p=0.006$). Differences in vessel morphology and white matter hyperintensity lesions between tertiles of the study group, divided based on the cardiovascular score, are presented in Figure 1.

Vessel Morphology and brain MRI biomarkers of cerebral blood flow, arrival time and white matter lesion count

To explore whether cerebral blood flow also varied with cardiovascular risk factors, a subgroup ($n=52$) analysis was performed in those with cerebral blood flow measures (mean cerebral blood flow 60 ml/100g/min (SD 11.5) and mean blood arrival time 1.01 seconds (SD 0.08). Slower blood arrival time (0.1 seconds per 1kg/m², 95%CI 0.001 to 0.05, $p=0.001$) and lower cerebral blood flow (-1.1 ml/100g/min per 1kg/m², 95%CI -2.0 to -0.1, $p=0.03$) were correlated with higher BMI (Supplementary Data, eTable 3). Cerebral blood flow was also lower in correlation with anti-hypertensive medication 11 ml/100g/min (95%CI -18 to -3, $p=0.007$). Cerebral blood flow was

2.5ml/100g/min higher for each additional healthier category of the cardiovascular score (95%CI 0.16 to 4.89, $p=0.03$). There was no significant correlation between blood arrival time and the cardiovascular scores (Table 4).

In multivariable analysis, controlling for modifiable risk factors (SBP, BMI, VPA, smoking, alcohol intake) blood arrival time and cerebral blood flow varied with cerebral vessel density, with each additional vessel per cm^3 correlating with a 0.015 seconds faster blood arrival time (95%CI -0.03 to -0.002, $p=0.02$) and 3 ml/100g/min increase in cerebral blood flow (95%CI 0.7 to 5.4, $p=0.01$). Vessel density was inversely correlated with white matter hyperintensity lesion count with 1.5 fewer lesions per unit increase in vessel density per cm^3 (95%CI to -2.7 to -0.4, $p=0.01$). (Table 3).

Discussion

In this cross-sectional study, optimal status of modifiable cardiovascular risk factors in young adults were associated with differences in brain vessel structure and function as well as a lower number of white matter hyperintensity lesions. Higher vessel density correlated with both higher cerebral blood flow and lower white matter lesion counts.

To date, studies tracking changes in brain vascular measures have largely focused on the transition from middle age to older adulthood. Cerebral blood flow is estimated to decline over the life course²⁶ with risk of dementia in older adults being 2 to 3 fold higher in those whose cerebral blood flow is below 55 ml/100g/min²⁷. Vascular dementia has also been associated with lower vascular density in brains of adults who have an early diagnoses of disease²⁸. In the current study, young adults in the lowest tertile for the modifiable cardiovascular score had approximately 1 vessel/ cm^3 lower vessel density and a mean value for cerebral blood flow of 55 ml/100g/min,

which is in the bottom 40% of the current study population. Therefore, the distribution of MRI findings observed in the current study raises the potential that some individuals may be starting to diverge on to different risk trajectories for brain vascular health in early adulthood. Furthermore, levels of cerebral blood flow associated with an increased risk of dementia are evident in some young adults. No participants had clinically significant white matter hyperintensity lesion volumes but lesion count was up to 4 lesions lower in the highest tertile of optimal status of modifiable risk factors.

Adverse modifiable cardiovascular risk factors are major determinants of white matter hyperintensity progression²⁹, with small lesions increasing in size or clustering into confluent lesions³⁰. Accumulation of lesions from an early age might explain why, by mid-life, white matter hyperintensity lesion volume is an established predictor of future stroke and dementia risk⁵. The longitudinal relationships between vessel morphology, cerebral perfusion and white matter lesion burden are uncertain. However, the patterns observed in the current study may suggest that resilience of the white matter, and potential to withstand risk exposures, may be influenced by the vascular morphology of an individual.

Modifiable risk factors such as blood pressure, BMI, smoking and lipid profile are known to drive systemic vascular disease in young adults in part through biological vascular disorders including endothelial dysfunction and oxidative stress³¹⁻³³. The current study suggests the cerebrovasculature may be similarly affected. Novel early life factors, such as preterm birth, are linked with early vascular disease³⁴ and the third trimester and early neonatal period are hypothesized to be times of significant vascular remodelling. In this study, gestational age was associated with vessel tortuosity, consistent with previous reports in infants³⁵, but not other cerebrovascular measures. Further work is needed to understand whether this was because

participants were largely born late preterm or because cardiovascular risk profile overwhelms this early exposure.²⁸.

The observed association between brain vascular measures and modifiable risk factors raises the potential for targeted intervention to prevent progression to disease. Reducing multiple risk factors can change risk trajectories and reduce vascular disease burden³⁶, with sustained lifestyle intervention and active blood pressure lowering associated with lower burden of white matter hyperintensity lesions and improved cerebral perfusion^{37, 38}. These interventions typically achieve 25% improvements in cardiovascular fitness and 10 mmHg reductions in blood pressure^{37, 38}, comparable to differences between high and low tertile groups for the cardiovascular scores in this study.

However, lifestyle-based primary cardiovascular prevention in young adults requires complex intervention design. Recent systematic review of interventions in young hypertensives demonstrated that the optimal way to intervene is poorly understood with lack of sustained effect³⁹. The alternative to lifestyle interventions would be pharmacological treatment. However, in this study group higher blood pressure was associated with reduced vessel density and anti-hypertensive use was associated with lower cerebral blood flow. Therefore, further work to identify optimal interventions in young adults to maintain autoregulation of cerebral blood flow, while reducing risk, may be required.

Limitations

This study has several limitations. First, a small sample recruited at a single site increases risk of bias and type 1 error while the study may be underpowered to identify subtle correlations with some risk factors. Second, purposive mixed passive and active recruitment strategies mean the sample is not population-based and could

be considered similar to a convenience sample. Therefore, it is not possible to generalise expected prevalence of cerebrovascular changes to the wider population. Third, the study is cross-sectional and causality or even temporality of the observed relationships cannot be inferred. Fourth, the cardiovascular risk assessment would be strengthened by detailed dietary questionnaires which were not included in this study. Fifth, cerebral blood flow was only available in a subgroup so ability to understand interactive effects of modifiable risk factors, vascular remodelling and perfusion on white matter integrity is limited. Sixth, longitudinal follow up will be required to determine the clinical significance of the observed findings. As such, this study should be considered preliminary and exploratory but does support a need for future work. The complexity of the imaging protocol and associated financial costs may limit its widespread use but large multi-centre studies with more focused protocols, and extended follow up, may have the potential to track vascular remodelling and assessment of impact on white matter and later disease.

Conclusion

In this preliminary study involving young adults without clinical evidence of cerebrovascular disease, modifiable cardiovascular risk factors were associated with MRI indices of cerebral vessel structure and function, and white matter hyperintensities. Further research is needed to verify these findings and determine clinical importance.

Acknowledgement

Study funding was provided by a British Heart Foundation (BHF) project grant (BHF Ref PG/13/58/30397), the Oxford BHF Centre for Research Excellence, and National Institute for Health Research (NIHR) Oxford Biomedical Research Centre. Dr Wilby Williamson is funded by a Wellcome Trust Clinical Research Training Fellowship (Ref 105741/Z/14/Z).

Authorship

All authors meet criteria for authorship: WW, AJL, HB, CF, HD, PL contributed to the design of the study, secured funding and refined the overall study protocol and lead the project delivery, NF, LG, TO, MJ, CM contributed to the development of the Brain MRI protocol and related pipelines, AJL, WW, OH, JF, SN contributed to image acquisition and quality control, WW, NF, LG, TO, MJ, CM, JB, HB, TS, DM, RP contributed to brain MRI image processing and analysis, AD advised on accelerometer protocol for objective physical activity measurement and completed analysis of raw data, WW, AJL, HB, OH, completed cardiovascular risk assessment and analysis of measures, WW, CF, AJL, PL and EF contributed to the statistical analysis, WW wrote the manuscript with support from LG, OH, AJL, CF, NF, HD, PL. All authors contributed to revision of the manuscript. PL completed the final edit of the manuscript.

Disclosures

Dr. Okell reports grants from The Royal Academy of Engineering, during the conduct of the study; In addition, Dr. Okell has a patent (US Patent 9,757,047) with royalties paid from Siemens Healthcare. All other authors declare no competing interests.

Role of the funding source

The funders of the study had no role in design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

Access to data

Dr. Williamson and Professor Leeson had full access to all of the data in the study and take full responsibility for the integrity of the data and the accuracy of the data analysis.

459 **References**

- 460 1. Raitakari OT, Juonala M, Kähönen M and et al. Cardiovascular risk factors in
461 childhood and carotid artery intima-media thickness in adulthood: The cardiovascular risk in
462 young finns study. *JAMA*. 2003;290:2277-2283.
- 463 2. Lewandowski AJ, Augustine D, Lamata P, Davis EF, Lazdam M, Francis J, McCormick
464 K, Wilkinson A, Singhal A, Lucas A, Smith N, Neubauer S and Leeson P. Preterm Heart in
465 Adult Life: Cardiovascular Magnetic Resonance Reveals Distinct Differences in Left
466 Ventricular Mass, Geometry and Function. *Circulation*. 2013;127:197-206.
- 467 3. Leeson CPM, Whincup PH, Cook DG, Donald AE, Papacosta O, Lucas A and Deanfield
468 JE. Flow-Mediated Dilation in 9- to 11-Year-Old Children. *Circulation*. 1997;96:2233.
- 469 4. Pase MP, Beiser A, Enserro D, Xanthakis V, Aparicio H, Satizabal CL, Himali JJ, Kase
470 CS, Vasan RS, DeCarli C and Seshadri S. Association of Ideal Cardiovascular Health With
471 Vascular Brain Injury and Incident Dementia. *Stroke*. 2016;47:1201-6.
- 472 5. Debette S and Markus HS. The clinical importance of white matter hyperintensities
473 on brain magnetic resonance imaging: systematic review and meta-analysis. *BMJ*.
474 2010;341:c3666.
- 475 6. van Dijk EJ, Prins ND, Vrooman HA, Hofman A, Koudstaal PJ and Breteler MM.
476 Progression of cerebral small vessel disease in relation to risk factors and cognitive
477 consequences: Rotterdam Scan study. *Stroke*. 2008;39:2712-9.
- 478 7. Hainsworth AH and Markus HS. Do in vivo experimental models reflect human
479 cerebral small vessel disease? A systematic review. *Journal of cerebral blood flow and*
480 *metabolism : official journal of the International Society of Cerebral Blood Flow and*
481 *Metabolism*. 2008;28:1877-91.
- 482 8. Yano Y, Stamler J, Garside DB, Daviglius ML, Franklin SS, Carnethon MR, Liu K,
483 Greenland P and Lloyd-Jones DM. Isolated systolic hypertension in young and middle-aged
484 adults and 31-year risk for cardiovascular mortality: the Chicago Heart Association Detection
485 Project in Industry study. *J Am Coll Cardiol*. 2015;65:327-35.
- 486 9. Aberg ND, Kuhn HG, Nyberg J, Waern M, Friberg P, Svensson J, Toren K, Rosengren
487 A, Aberg MA and Nilsson M. Influence of Cardiovascular Fitness and Muscle Strength in Early
488 Adulthood on Long-Term Risk of Stroke in Swedish Men. *Stroke*. 2015;46:1769-76.
- 489 10. Hartiala O, Kajander S, Knuuti J, Ukkonen H, Saraste A, Rinta-Kiikka I, Kainulainen S,
490 Kahonen M, Hutri-Kahonen N, Laitinen T, Lehtimäki T, Viikari JS, Hartiala J, Juonala M,
491 Raitakari OT and Magnussen CG. Life-course risk factor levels and coronary artery
492 calcification. The Cardiovascular Risk in Young Finns Study. *Int J Cardiol*. 2016;225:23-29.
- 493 11. Griffanti L, Zamboni G, Khan A, Li L, Bonifacio G, Sundaresan V, Schulz UG, Kuker W,
494 Battaglini M, Rothwell PM and Jenkinson M. BIANCA (Brain Intensity AbNormality
495 Classification Algorithm): A new tool for automated segmentation of white matter
496 hyperintensities. *NeuroImage*. 2016;141:191-205.
- 497 12. Forkert ND, Schmidt-Richberg A, Fiehler J, Illies T, Möller D, Handels H and Säring D.
498 Fuzzy-based Vascular Structure Enhancement in Time-of-Flight MRA Images for Improved
499 Segmentation. *Methods of Information in Medicine*. 2011;50:74-83.
- 500 13. Okell TW, Chappell MA, Kelly ME and Jezzard P. Cerebral Blood Flow Quantification
501 Using Vessel-Encoded Arterial Spin Labeling. *Journal of Cerebral Blood Flow & Metabolism*.
502 2013;33:1716-1724.
- 503 14. Berry ES, Jezzard P and Okell TW. An Optimized Encoding Scheme for Planning
504 Vessel-Encoded Pseudocontinuous Arterial Spin Labeling. *Magn Reson Med*. 2015;74:1248-
505 56.
- 506 15. Chappell MA, Okell TW, Payne SJ, Jezzard P and Woolrich MW. A fast analysis
507 method for non-invasive imaging of blood flow in individual cerebral arteries using vessel-
508 encoded arterial spin labelling angiography. *Medical Image Analysis*. 2012;16:831-839.

16. Forkert ND, Schmidt-Richberg A, Fiehler J, Illies T, Möller D, Handels H and Säring D. Automatic Correction of Gaps in Cerebrovascular Segmentations Extracted from 3D Time-of-Flight MRA Datasets. *Methods of Information in Medicine*. 2012;51:415-422.
17. Foster CE, Brennan G, Matthews A, McAdam C, Fitzsimons C and Mutrie N. Recruiting participants to walking intervention studies: a systematic review. *Int J Behav Nutr Phys Act*. 2011;8:137.
18. Aigner A, Grittner U, Rolfs A, Norrving B, Siegerink B and Busch MA. Contribution of Established Stroke Risk Factors to the Burden of Stroke in Young Adults. *Stroke*. 2017;48:1744-1751.
19. Seshadri S, Wolf PA, Beiser A, Elias MF, Au R, Kase CS, D'Agostino RB and DeCarli C. Stroke risk profile, brain volume, and cognitive function: The Framingham Offspring Study. *Neurology*. 2004;63:1591-1599.
20. Spartano. Midlife exercise blood pressure, heart rate, and fitness relate to brain volume 2 decades later. *Neurology*. 2016;86:1313-1319.
21. Smith SM, Jenkinson M, Woolrich MW, Beckmann CF, Behrens TE, Johansen-Berg H, Bannister PR, De Luca M, Drobnjak I, Flitney DE, Niazy RK, Saunders J, Vickers J, Zhang Y, De Stefano N, Brady JM and Matthews PM. Advances in functional and structural MR image analysis and implementation as FSL. *NeuroImage*. 2004;23 Suppl 1:S208-19.
22. Siepmann T, Boardman H, Bilderbeck A, Griffanti L, Kenworthy Y, Zwager C, McKean D, Francis J, Neubauer S, Yu GZ, Lewandowski AJ, Sverrisdottir YB and Leeson P. Long-term cerebral white and gray matter changes after preeclampsia. *Neurology*. 2017;88:1256-1264.
23. Sexton CE, Zsoldos E, Filippini N, Griffanti L, Winkler A, Mahmood A, Allan CL, Topiwala A, Kyle SD, Spiegelhalter K, Singh-Manoux A, Kivimaki M, Mackay CE, Johansen-Berg H and Ebmeier KP. Associations between self-reported sleep quality and white matter in community-dwelling older adults: A prospective cohort study. *Hum Brain Mapp*. 2017;38:5465-5473.
24. Hopkins RO, Beck CJ, Burnett DL, Weaver LK, Victoroff J and Bigler ED. Prevalence of white matter hyperintensities in a young healthy population. *J Neuroimaging*. 2006;16:243-51.
25. Mersov AM, Crane DE, Chappell MA, Black SE and MacIntosh BJ. Estimating the sample size required to detect an arterial spin labelling magnetic resonance imaging perfusion abnormality in voxel-wise group analyses. *Journal of Neuroscience Methods*. 2015;245:169-177.
26. Chen JJ, Rosas HD and Salat DH. Age-associated reductions in cerebral blood flow are independent from regional atrophy. *NeuroImage*. 2011;55:468-78.
27. Wolters FJ, Zonneveld HI, Hofman A, van der Lugt A, Koudstaal PJ, Vernooij MW, Ikram MA and Heart-Brain Connection Collaborative Research G. Cerebral Perfusion and the Risk of Dementia: A Population-Based Study. *Circulation*. 2017;136:719-728.
28. Brown WR, Moody DM, Thore CR, Challa VR and Anstrom JA. Vascular dementia in leukoaraiosis may be a consequence of capillary loss not only in the lesions, but in normal-appearing white matter and cortex as well. *J Neurol Sci*. 2007;257:62-6.
29. Dufouil C, Kersaint-Gilly A, Besançon V, Levy C, Auffray E and Brunnereau L. Longitudinal study of blood pressure and white matter hyperintensities: the EVA MRI Cohort. *Neurology*. 2001;56.
30. Schmidt R, Enzinger C, Ropele S, Schmidt H and Fazekas F. Progression of cerebral white matter lesions: 6-year results of the Austrian Stroke Prevention Study. *The Lancet*. 361:2046-2048.
31. Berenson GS, Srinivasan SR, Bao W, Newman WP, Tracy RE and Wattigney WA. Association between Multiple Cardiovascular Risk Factors and Atherosclerosis in Children and Young Adults. *New England Journal of Medicine*. 1998;338:1650-1656.

32. Stamler J, Daviglus ML, Garside DB, Dyer AR, Greenland P and Neaton JD. Relationship of baseline serum cholesterol levels in 3 large cohorts of younger men to long-term coronary, cardiovascular, and all-cause mortality and to longevity. *JAMA*. 2000;284:311-318.
33. Relationship of atherosclerosis in young men to serum lipoprotein cholesterol concentrations and smoking: A preliminary report from the pathobiological determinants of atherosclerosis in youth (pday) research group. *JAMA*. 1990;264:3018-3024.
34. Raikonen K, Kajantie E, Pesonen AK, Heinonen K, Alastalo H, Leskinen JT, Nyman K, Henriksson M, Lahti J, Lahti M, Pyhala R, Tuovinen S, Osmond C, Barker DJ and Eriksson JG. Early life origins cognitive decline: findings in elderly men in the Helsinki Birth Cohort Study. *PLoS one*. 2013;8:e54707.
35. Vasu V, Durighel G, Thomas EL, Malamateniou C, Bell JD, Rutherford MA and Modi N. Preterm nutritional intake and MRI phenotype at term age: a prospective observational study. *BMJ open*. 2014;4:e005390.
36. Spring B, Moller AC, Colangelo LA, Siddique J, Roehrig M, Daviglus ML, Polak JF, Reis JP, Sidney S and Liu K. Healthy lifestyle change and subclinical atherosclerosis in young adults: Coronary Artery Risk Development in Young Adults (CARDIA) study. *Circulation*. 2014;130:10-7.
37. Espeland MA, Erickson K, Neiberg RH, Jakicic JM, Wadden TA, Wing RR, Desiderio L, Erus G, Hsieh M-K, Davatzikos C, Maschak-Carey BJ, Laurienti PJ, Demos-McDermott K and Bryan RN. Brain and White Matter Hyperintensity Volumes After 10 Years of Random Assignment to Lifestyle Intervention. *Diabetes Care*. 2016;39:764.
38. Espeland MA, Luchsinger JA, Neiberg RH, Carmichael O, Laurienti PJ, Pi-Sunyer X, Wing RR, Cook D, Horton E, Casanova R, Erickson K, Nick Bryan R and Action for Health in Diabetes Brain Magnetic Resonance Imaging Research G. Long Term Effect of Intensive Lifestyle Intervention on Cerebral Blood Flow. *J Am Geriatr Soc*. 2018;66:120-126.
39. Williamson W, Foster C, Reid H, Kelly P, Lewandowski AJ, Boardman H, Roberts N, McCartney D, Huckstep O, Newton J, Dawes H, Gerry S and Leeson P. Will Exercise Advice Be Sufficient for Treatment of Young Adults With Prehypertension and Hypertension? A Systematic Review and Meta-Analysis. *Hypertension*. 2016;68:78-87.

591 Table 1. **Age, demographics and cardiovascular risk profile of study group.**

	Study Group (n=125)
Demographics	
Age, mean (SD), years	24.7 (5.0)
Female, n, (%)	61 (49%)
Gestational Age, mean (SD), weeks	36.6 (4.3)
Smoking, n, (%)	19 (15.2)
Smokers' median pack years (Q1-Q3)	2.7 (6.7)
Alcohol, n, (%)	97 (77.6)
Alcohol consumers' median drinks per week (Q1-Q3)	4.0 (4.0)
Hypertension Diagnosis, n, (%)	29 (23.0)
Taking Hypertension Medication, n (%)	21 (16.8)
FHx Stroke or CHD, n, (%)	10 (8)
Education Level	
Completed University, n, (%)	86 (68.8)
Anthropometrics	
Height, mean (SD), m	1.73 (0.1)
Weight, mean (SD), kg	70.9 (13.8)
BMI, mean (SD), kg/m ²	23.6 (3.7)
Blood pressure, mean (SD), mmHg	
Resting Systolic	122.0 (11.6)
Resting Diastolic	71.3 (9.55)
Ambulatory Awake Systolic	129.6 (11.8)
Ambulatory Awake Diastolic	76.9 (8.0)
Peak Exercise Systolic	174.8 (25.4)
Peak Exercise Diastolic	87.1 (12.4)
Fitness	
Peak VO ₂ , mean (SD), ml/kg/min	37.9 (9.6)
Peak Respiratory Exchange Ratio, mean (SD)	1.2 (0.06)
VPA, median (Q1-Q3), hours per week	0.74 (1.25)
MVPA, median (Q1-Q3), hours per week	14.73 (6.09)
Biochemistry	
Total Cholesterol, mean (SD), mg/dL	170.15 (29.0)
LDL, mean (SD), mg/dL	97.45 (25.9)
HDL, mean (SD), mg/dL	55.68 (11.2)
TChol:HDL ratio, mean (SD)	3.18 (0.85)
Triglyceride, median (IQR), mg/dL	74.4 (54.0)
Blood Glucose, mean (SD), mg/dL	88.2 (7)
HOMA-IR, mean (SD)	0.77 (0.46)
HsCRP, median (Q1-Q3), mg/L	0.57 (1.16)
Brain MRI Vessel, Perfusion and White Matter Parameters	
Brain vessel density, mean (SD), vessels/cm ³	8.3 (1.41)
Brain vessel calibre, mean (SD), µm	531 (36)
Brain vessel tortuosity, mean (SD)	1.49 (0.088)
Brain white matter hyperintensity lesion count, mean (SD)	20.9 (7.9)
Brain Blood Arrival Time (SD), seconds	1.01 (0.08)
Cerebral Blood Flow (SD), ml/100g/min	60 (11.5)

592 Abbreviations: FHx, Family History, BMI, body mass index; SBP, systolic blood
593 pressure; DBP, diastolic blood pressure; Peak VO₂, Peak Oxygen Uptake; VPA,
594 Vigorous Physical Activity; MVPA, Moderate to Vigorous Physical Activity; LDL, low
595 density lipoprotein; HDL, high density lipoprotein; T Chol: total cholesterol; HsCRP,
596 highly sensitive C reactive protein; HOMA-IR, homeostatic model assessment of

597 insulin resistance. Brain blood arrival time and cerebral blood flow data was available
598 in 52 participants.
599

Table 2. Association between non-modifiable and modifiable risk factors and brain vessel morphology (vessel density, caliber and tortuosity).

	Bivariable Point Estimate (95 %CI)	P value	Adjusted Point Estimate (95 %CI)	P value
Brain Vessel Density (vessels/cm³)			Model Statistics R²=0.20 p =.009	
Gestational Age, weeks	-0.001 (-0.06 to 0.06)	.98	-0.02 (-0.08 to 0.03)	.42
Resting SBP, mmHg	-0.03 (-0.05 to -0.004)	.02	-0.02 (-0.04 to -0.0004)	.046
BMI, kg/m ²	-0.10 (-0.16 to -0.02)	.01	-0.08 (-0.15 to -0.01)	.02
VPA, hours per week	0.10 (-0.17 to 0.39)	.42	-0.04 (-0.28 to 0.20)	.75
Alcoholic drinks per week	-0.10 (-0.025 to -0.008)	.31	-0.01 (-0.04 to 0.02)	.41
Smoking pack years	0.20 (0.06 to 0.30)	.004	0.17(0.06 to 0.28)	.004
Peak VO ₂ , ml/kg/min	0.01 (-0.02 to 0.04)	.5	.	.
Peak Ex DBP, mmHg	-0.02 (-0.04 to -0.003)	.047	.	.
Cholesterol/HDL Ratio	-0.40 (-0.69 to -0.06)	.02	.	.
HOMA IR	-0.56 (-1.17 to 0.04)	.07	.	.
Hypertension Rx	0.75 (-0.01 to 1.5)	.05	.	.
Brain Vessel Caliber (μm)			Model Statistics R²=0.24 p=.001	
Gestational Age, weeks	-0.1 (-2.0 to 1.0)	.88	-1.0 (-3.0 to 0.5)	.16
Resting SBP, mmHg	-0.4 (-1.0 to 2.0)	.15	-0.6 (-1.0 to -0.05)	.03
BMI, kg/m ²	-1.0 (-3.0 to 1.0)	.33	-1.0 (-3.0 to 1.0)	.42
VPA, hours per week	1.0 (-6.0 to 8.0)	.73	-2.0 (-9.0 to 4.0)	.49
Alcoholic drinks per week	-0.1 (-1.0 to 1.0)	.70	-1.0 (-2.0 to 0.1)	.09
Smoking pack years	3.0 (-0.2 to 6.0)	.06	4.0 (0.2 to 8.0)	.04
Peak VO ₂ , ml/kg/min	0.4 (-0.2 to 1.0)	.19	.	.
Peak Ex DBP, mmHg	-1.0 (-1.4 to -0.4)	<.001	.	.
Cholesterol/HDL Ratio	-3.0 (-10.0 to 5.0)	.52	.	.
HOMA IR	-14.0 (-30 to 1.0)	.08	.	.
Hypertension Rx	10 (-9.0 to 31.0)	.27	.	.
Brain Vessel Tortuosity			Model Statistics R²=0.1 p=.26	
Gestational Age, weeks	0.005 (0.001 to 0.009)	.007	0.006 (0.001 to 0.01)	.01

Also modelled was the association between these risk factors and tortuosity, and only gestational age was related, full analysis presented in supplement (eTable 1). Model adjusted for non-modifiable factors of age, sex, gestational age and modifiable risk factors of systolic blood pressure, body mass index, vigorous physical activity, weekly alcohol consumption and smoking status. Abbreviations and units: SBP, systolic blood pressure (mmHg); BMI, body mass index (kg/m²); VPA, Vigorous Physical Activity (hours per week); Alcohol (drinks per week); Smoking (pack years); Peak VO₂, Peak Oxygen Uptake (ml/kg/min); Ex DBP, Peak exercise diastolic blood pressure (mmHg), Cholesterol/HDL ratio, ratio total cholesterol/high density lipoprotein; HOMA-IR, homeostatic model assessment of insulin resistance,

613 Hypertension Rx participant taking prescription medications for hypertension
614 (yes/no). Exposure variables were available for all participants. The point estimate
615 refers to the magnitude of change in the vessel morphology variable per unit change
616 in the non-modifiable and modifiable variables.

617

Table 3. Association of vessel morphology (density, caliber, tortuosity) with measures of brain blood arrival time, cerebral blood flow and white matter hyper-intensity lesions

	Bivariable Point Estimate (95 %CI)	P Value	Adjusted Point Estimate (95 %CI)	P Value
Blood Arrival Time (seconds) (n=52)				
Brain Vessel Density, vessels/cm ³	-0.03 (-0.04 to -0.01)	.002	-0.015 (-0.03 to -0.002)	.02
Brain Vessel Caliber, µm	0.08 (-0.61 to 0.78)	.81	0.22 (-0.28 to 0.71)	.38
Brain Vessel Tortuosity	0.13 (-0.15 to 0.4)	0.36	-0.014 (-0.23 to 0.21)	.90
Cerebral Blood Flow (ml/100g/min) (n=52)				
Brain Vessel Density, vessels/cm ³	4.0 (1.8 to 6.2)	.001	3.1 (0.7 to 5.4)	.01
Brain Vessel Caliber, µm	48.6 (-50.3 to 147.6)	.34	-8.0 (-126.1 to 110.1)	.89
Brain Vessel Tortuosity	3.8 (-36.4 to 44.1)	.85	12.9 (-35.4 to 61.1)	.60
White Matter Hyperintensity Lesion Count (lesions) (n=125)				
Brain Vessel Density, vessels/cm ³	-1.1 (-2.2 to 0.06)	.06	-1.5 (-2.7 to -0.4)	.01
Brain Vessel Caliber, µm	13.5 (-31.3 to 58.4)	.55	12.1 (-34.5 to 57.8)	.61
Brain Vessel Tortuosity	-17.5(-35.3 to 0.24)	.05	-11.0 (-29.0 to 7.0)	.23

Model adjusted for non-modifiable factors of age, sex, gestational age and modifiable risk factors of systolic blood pressure, body mass index, vigorous physical activity, weekly alcohol consumption and smoking status. Also modelled were the association with vessel caliber and tortuosity, only vessel density was related. The point estimate refers to the magnitude of change in blood arrival time, cerebral blood flow or number of white matter hyperintensity lesions per unit change in respective vessel morphological variable.

Table 4. Modifiable cardiovascular score and association with brain vessel morphology, cerebral blood flow and white matter hyperintensity lesion count


Modifiable Cardiovascular Score	Brain Vessel Density, vessels/cm ³ point estimate (95%CI)	Brain Vessel Caliber, μ m point estimate (95%CI)	Brain Vessel Tortuosity point estimate (95%CI)	Brain Blood Arrival Time, seconds (n=52) point estimate (95%CI)	Cerebral Blood Flow, ml/min/100g (n=52) point estimate (95%CI)	Brain white matter hyperintensity lesion count, number point estimate (95%CI)
1 (n=2)	9.2 (6.4 to 11.9)	505 (437 to 573)	1.52 (1.44 to 1.50)	.	.	36.2 (24.3 to 48.0)
2 (n=0)
3 (n=4)	6.9 (5.0 to 8.8)	518 (470 to 565)	1.49 (1.46 to 1.52)	1.26 (1.16 to 1.36)	66.6 (50.4 to 82.0)	24.0 (16 to 32)
4 (n=14)	7.4 (6.6 to 8.2)	512 (493 to 532)	1.47 (1.45 to 1.53)	1.22 (1.16 to 1.27)	54.2 (45.5 to 63.0)	25.0 (21.0 to 29.2)
5 (n=27)	8.0 (7.4 to 8.5)	524 (510 to 540)	1.51 (1.47 to 1.55)	1.21 (1.16 to 1.26)	54.6 (47.0 to 62.0)	22 (19.0 to 25.3)
6 (n=36)	8.5 (8.0 to 9.0)	533 (521 to 545)	1.49 (1.46 to 1.52)	1.19 (1.15 to 1.23)	60.2 (54.0 to 67.0)	21.0 (19.0 to 24.0)
7 (n=33)	8.5 (8.0 to 9.0)	542 (530 to 555)	1.48 (1.45 to 1.52)	1.18 (1.14 to 1.22)	64.0 (57.8 to 70.0)	19.0 (16.2 to 21.8)
8 (n=9)	9.1 (8.2 to 10.0)	540 (518 to 563)	1.54 (1.46 to 1.62)	1.18 (1.11 to 1.24)	68.0 (57.6 to 78.1)	20.0 (15.4 to 26.6)
Change in point estimate per additional score (n=125)	0.31 (0.112 to 0.514)	8.0 (3.0 to 13.0)	0.005 (-0.008 to 0.18)	-0.014 (-0.03 to 0.001)	2.5 (0.16 to 4.89)	-1.6 (-3.0 to -0.5)

Participants were assessed for a cardiovascular score, for each healthier category of a modifiable risk factor according to the following criteria: BMI <25 kg/m²; highest tertile cardiovascular fitness and/or physical activity; alcohol <8 drinks/week; non-smoker for >6 months; blood pressure on awake ambulatory monitoring <130/80 mmHg; a non-hypertensive diastolic response to exercise (peak diastolic blood pressure <90 mmHg), total cholesterol <200mg/dL; and fasting glucose <100mg/dL. Adjusted for age and sex. The point estimate refers to the magnitude of change in the dependent variable per unit change in the modifiable cardiovascular score.

Figure 1. Comparison of white matter lesion count and vessel morphology between groups of participants based on their modifiable cardiovascular score.

The cardiovascular score provided a cumulative score for each of the following factors: high cardiovascular fitness (top tertile of peak oxygen uptake ($\geq 110\%$ predicted peak oxygen uptake) or participating in ≥ 75 minutes vigorous physical activity per week); not smoking in last 6 months; alcohol < 8 drinks/week; ambulatory awake blood pressure $< 130/80$ mmHg; body mass index < 25 kg/m²; fasting total cholesterol < 200 mg/d; fasting blood glucose < 100 mg/dL; and diastolic blood pressure at peak exercise ≤ 90 mmHg. Figure 1 presents a post-hoc comparisons between groups of participants who score 0 to 5 positive factors (n=47), 6 factors (n=36) and 7 to 8 positive factors (n=42). The groupings were defined to approximate tertiles of the combined cardiovascular score. Panel A presents the white matter lesions counts for individual participants and associated group mean and 95% CI, Panels B and C present the mean group values and 95% CI. Participants with 7 to 8 healthier categories of risk factor have a mean vessel density 11% higher than participants with 0 to 5 healthier categories of risk factor (Panel B, 8.6 vessels/cm³ (SD 1.39) vs 7.8 vessels/cm³ (SD 1.21) p=0.007), a mean vessel caliber 3% higher (Panel C, 538 μ m (SD 21) vs 522 μ m (SD 45) p=0.02) and on average 20% lower white matter hyperintensity lesion counts (Panel A, 19.6 lesions (SD 7.8) vs 23.5 lesions (SD 8.6) p=0.03). Panels present group means and 95%CI and reported group differences are adjusted for age and sex.


Figure 1. Comparison of white matter lesion count and vessel morphology between groups of participants based on their modifiable cardiovascular score.




The cardiovascular score provided a cumulative score for each of the following factors: high cardiovascular fitness (top tertile of peak oxygen uptake ($\geq 110\%$ predicted peak oxygen uptake) or participating in ≥ 75 minutes vigorous physical activity per week); not smoking in last 6 months; alcohol < 8 drinks/week; ambulatory awake blood pressure $< 130/80$ mmHg; body mass index < 25 kg/m²; fasting total cholesterol < 200 mg/d; fasting blood glucose < 100 mg/dL; and diastolic blood pressure at peak exercise ≤ 90 mmHg. Figure 1 presents a post-hoc comparisons between groups of participants who score 0 to 5 positive factors (n=47), 6 factors (n=36) and 7 to 8 positive factors (n=42). The groupings were defined to approximate tertiles of the combined cardiovascular score. Panel A presents the white matter lesions counts for individual participants and associated group mean and 95% CI, Panels B and C present the mean group values and 95% CI. Participants with 7 to 8 healthier categories of risk factor have a mean vessel density 11% higher than participants with 0 to 5 healthier categories of risk factor (Panel B, 8.6 vessels/cm³ (SD 1.39) vs 7.8 vessels/cm³ (SD 1.21) $p=0.007$), a mean vessel caliber 3% higher (Panel C, 538 μ m (SD 21) vs 522 μ m (SD 45) $p=0.02$) and on average 20% lower white matter hyperintensity lesion counts (Panel A, 19.6 lesions (SD 7.8) vs 23.5 lesions (SD 8.6) $p=0.03$). Panels present group means and 95%CI and reported group differences are adjusted for age and sex.

**White Matter Lesion Count
(lesions/total white matter)**

Panel A



Panel B



Panel C

